

## Description of *Clostridium phoceensis* sp. nov., a new species within the genus *Clostridium*

M. Hosny, S. Benamar, G. Durand, N. Armstrong, C. Michelle, F. Cadoret, B. La Scola and N. Cassir

Unité de Recherche sur les Maladies Infectieuses et Tropicales Emergentes (URMITE), UM63 CNRS 7278 IRD 198 INSERM U1095, IHU Méditerranée Infection, Pôle des Maladies Infectieuses, Assistance Publique-Hôpitaux de Marseille, Faculté de Médecine, Marseille, France

### Abstract

*Clostridium phoceensis* sp. nov., strain GD3<sup>T</sup> (= CSUR P1929 = DSM 100334) is the type strain of *C. phoceensis* sp. nov., a new species within the genus *Clostridium*. This strain was isolated from the gut microbiota of a 28-year-old healthy French man. *C. phoceensis* is a Gram-negative, spore-forming, nonmotile, strictly anaerobic bacterium. We describe its complete genome sequence and annotation, together with its phenotypic characteristics.

© 2016 The Authors. Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases.

**Keywords:** Anaerobe, *Clostridium phoceensis*, culturomics, gut microbiota, taxono-genomics

**Original Submission:** 17 June 2016; **Revised Submission:** 9 September 2016; **Accepted:** 16 September 2016

**Article published online:** 25 September 2016

**Corresponding author:** N. Cassir, Unité de Recherche sur les Maladies Infectieuses et Tropicales Emergentes, URMITE, UM63, CNRS 7278, IRD 198, INSERM 1095, Faculté de médecine, Aix-Marseille Université, 27 Boulevard Jean Moulin, 13385 Marseille Cedex 05, France

**E-mail:** [cassirnadim@gmail.com](mailto:cassirnadim@gmail.com)

M. Hosny and S. Benamar contributed equally to this article, and both should be considered first author.

### Introduction

Human adult gut microbiota is estimated to consist of up to 100 trillion microorganisms, comprising at least 500 different species, mostly anaerobic bacteria [1]. Although the advent of modern molecular microbiological methods has expanded the degree of bacterial detection from stool samples, it does not allow for the phenotypic description of new living species [2]. Consequently, there has been renewed interest in culture methods [3].

A new taxonomic approach known as taxonogenomics has recently been proposed to describe new isolated bacterial species [4]. This polyphasic strategy combines phenotypic characteristics, the matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) spectrum,

and the analysis and comparison of the complete genome sequence.

Since the creation of the genus *Clostridium* in 1880, more than 200 species have been described [5]. While species belonging to the family *Clostridiaceae* are mainly associated with the commensal digestive flora of mammals and can be commonly found in the environment, some are major human pathogens, including toxigenic *Clostridium botulinum*, *Peptoclostridium difficile*, *Clostridium tetani* and *Clostridium perfringens* [6].

We propose *Clostridium phoceensis* sp. nov., strain GD3<sup>T</sup> (= CSUR P1929 = DSM 100334) as the type strain of *C. phoceensis* sp. nov., a new species within the genus *Clostridium*. This strain was isolated from the gut microbiota of a 28-year-old healthy French man as part of a culturomics study aiming at individually cultivating all bacterial species from a stool sample. Here we describe the characteristics of *C. phoceensis* sp. nov. strain GD3<sup>T</sup>, including its phenotype and genome sequence.

### Materials and Methods

#### Sample collection

The stool sample was taken from a 28-year-old healthy French man. The sample was collected as part of a research study on the human gut microbiota. The study was approved by the

Institut Fédératif de Recherche 48 (agreement no. 09-022, Marseille, France), and the patient's consent was obtained. The sample was stored at  $-80^{\circ}\text{C}$  in La Timone Hospital (Marseille, France).

### Strain isolation and identification (MALDI-TOF MS and 16S rRNA sequencing)

The faecal sample was treated using the concept of culturomics [7]. The colonies obtained were identified using MALDI-TOF MS [8,9] (Bruker Daltonics, Leipzig, Germany) and analysed using a Microflex spectrometer (Bruker), leading to the protein spectrum being obtained. A score under 1.7 did not enable any identification. Subsequently, 16S rRNA was sequenced and the sequence was matched using BLAST (Basic Local Alignment Search Tool) against the National Center for Biotechnology Information database [10]. DNA was extracted using the EZ1 Tissue Extraction Kit (Qiagen, Hilden, Germany), and sequences were aligned using ChromasPro 1.6 (Technelysium, South Brisbane, Queensland, Australia).

### Growth conditions

The growth condition of the strain was determined by testing different temperatures and atmospheres. Five growth temperatures (ambient, 28, 37, 45 and  $56^{\circ}\text{C}$ ) were tested under anaerobic (GENbag anaer) microaerophilic atmospheres (GENbag microer) (bioMérieux, Marcy l'Étoile, France) and aerobic condition on 5% sheep's blood agar (bioMérieux). Colonies were obtained after thermal shock for 20 minutes at  $80^{\circ}\text{C}$  in an anaerobic blood culture bottle (Bactec Lytic/10 Anaerobic/F) supplemented with 5% sheep's blood at  $37^{\circ}\text{C}$ .

### Phenotypic, biochemical and antibiotic susceptibility tests

Gram staining, motility, catalase and oxidase were determined as described by Lagier *et al.* [3]. Sporulation was tested using a thermal shock on bacterial colonies (diluted in phosphate-buffered saline) for 10 minutes at  $80^{\circ}\text{C}$ . The biochemical characteristics were tested using API 50CH, API ZYM and API 20A strips (bioMérieux). Antibiotic susceptibility referred to European Committee on Antimicrobial Susceptibility Testing 2015 recommendations.

### Fatty acid methyl ester (FAME) analysis by gas chromatography/mass spectrometry (GC/MS)

FAME analysis was performed by GC/MS. Two samples were prepared with 2 mg of bacterial biomass from several culture plates. Two samples were prepared with approximately 2 mg of bacterial biomass per tube collected from several culture plates. FAMES were prepared as described by Sasser ([http://www.midi-inc.com/pdf/MIS\\_Technote\\_101.pdf](http://www.midi-inc.com/pdf/MIS_Technote_101.pdf)). GC/MS analyses were

carried out as described by Dione *et al.* [11]. Briefly, FAMES were separated using an Elite 5-MS column and monitored by mass spectrometry (Clarus 500-SQ8S; PerkinElmer, Courtaboeuf, France). A spectral database search was performed using MS Search 2.0 operated using the Standard Reference Database 1A (National Institute of Standards and Technology, Gaithersburg, MD, USA) and the FAME mass spectral database (Wiley, Chichester, UK).

### Microscopy


Individual cells of *C. phoceensis* strain GD3<sup>T</sup> were captured using a Tecnai G20 electron microscopy (FEI Company, Limeil-Brevannes, France). A Color Gram 2 Kit (bioMérieux) was used to perform the Gram coloration observed with a  $100\times$  oil-immersion objective lens using the DM1000 photonic microscope (Leica, Wetzlar, Germany) [12,13].

### Genome project history

The organism was selected for sequencing because it had been isolated from a healthy person for the first time and on the basis of its 16S rRNA similarity, phylogenetic position and phenotypic differences with other members of the genus *Clostridia*. The GenBank accession number is [CVUG01000000](https://www.ncbi.nlm.nih.gov/nucl/1000000) and consists of 10 scaffolds with a total of 16 contigs [14].

### Genome sequencing and assembly

The genomic DNA of *Clostridium phoceensis* GD3<sup>T</sup> was sequenced on the MiSeq sequencer (Illumina, San Diego, CA, USA) using the mate-pair strategy. The gDNA was barcoded in order to be mixed with 11 other projects using the Nextera mate-pair sample prep kit (Illumina). The mate-pair library was prepared with 1.5  $\mu\text{g}$  of genomic DNA using the Nextera Mate-Pair Illumina guide. The gDNA sample was simultaneously fragmented and tagged using a mate-pair junction adapter. The pattern of the fragmentation was validated on an Agilent 2100 BioAnalyzer (Agilent Technologies, Santa Clara, CA, USA) with a DNA 7500 lab chip. The DNA fragments ranged in size from 1 to 10 kb, with an optimal size of 4.490 kb. No size selection was performed, and only 600 ng of tagmented fragments were circularized. The circularized DNA was mechanically sheared to small fragments with an optimal size of 938 bp on the Covaris S2 device in microtubes (Covaris, Woburn, MA, USA). The library profile was visualized on a High Sensitivity Bioanalyzer LabChip (Agilent Technologies), and the final library concentration was measured at 4.457 nmol/L. The libraries were normalized at 2 nM and pooled. After a denaturation step and dilution at 15 pM, the pool of libraries was loaded onto the reagent cartridge and then onto the instrument along with the flow cell. Automated cluster generation and sequencing run were performed in a single 39-hour run in a  $2 \times 251$  bp; 6.1 Gb



**FIG. 1.** Reference mass spectrum (matrix-assisted desorption ionization–time of flight mass spectrometry) from *Clostridium phoceensis* strain GD3<sup>T</sup>.

of total information was obtained from a 653K/mm<sup>2</sup> cluster density, with a cluster passing quality control filters of 96.1% (12 031 000 clusters). Within this run, the index representation for *C. phoceensis* GD3<sup>T</sup> was determined to 9.32%. The 1 121 200 paired reads were filtered according to the read qualities. These reads were trimmed, then assembled by SPDES software. Finally, the draft genome of *C. phoceensis* GD3<sup>T</sup> consisted of ten scaffolds with 16 contigs and generated a genome size of 3.4 Mb with 59.32% G+C content.


#### Genome annotation

Open reading frames (ORFs) were predicted using Prodigal [15] with default parameters, although the predicted ORFs were excluded if they spanned a sequencing gap region. The predicted bacterial protein sequences were searched against the GenBank database [16] and the Clusters of Orthologous Groups (COGs) databases using BLASTP. The tRNA genes were found using the tRNAScanSE tool [17], while RNAmmer [18] was used to find ribosomal RNAs and BLASTn against the GenBank database. Lipoprotein signal peptides and the number of transmembrane helices were predicted using SignalP [19] and TMHMM [20] respectively. Artemis [21] was used for data management, and DNA Plotter [22] was used to visualize genomic features. To estimate the mean level of nucleotide sequence similarity at the genome level, we used homemade average genomic identity of orthologous gene sequences (AGIOS) software [4].

## Results

#### Strain isolation, identification and phylogeny

MALDI-TOF MS failed to identify the strain GD3<sup>T</sup>, so its mass spectrum was added to the Bruker database (Fig. 1). To improve identification, 16S rRNA sequencing was performed, and the access number in 16S rRNA EMBL-EBI (European Molecular Biology Laboratory–European Bioinformatics Institute) was assigned as LN846907. The highest value of nucleotide sequence similarity was observed with *Flavonifractor plautii* (97%), the phylogenetically closest species.



**FIG. 2.** Gram staining of *Clostridium phoceensis* strain GD3<sup>T</sup>.

### Phenotypic and biochemical characterization

*C. phoceensis* strain GD3<sup>T</sup> is a Gram-negative (Fig. 2), spore-forming, nonmotile, strictly anaerobic bacterium that has no catalase and oxidase activities, measuring 1.8 µm in length and 0.5 µm in diameter (Fig. 3). The sporulation test was positive; the organism grows at 45°C in anaerobic conditions. Using the API ZYM gallery, *C. phoceensis* exhibits esterase (C4), phosphatase acid, naphthol-AS-BI-phosphohydrolase and β-glucosidase activities. When using the API 20A gallery, a positive reaction was only observed for D-glucose (Table 1).

Of the antibiotics tested, *C. phoceensis* was found to be sensitive to amoxicillin, amoxicillin–clavulanate, ceftriaxone, ceftazidime, imipenem, doripenem and ciprofloxacin.

### Predominant cellular fatty acids

The predominant cellular fatty acids of *C. phoceensis* strain GD3<sup>T</sup> are hexadecanoic acid (16:0; 33.1 ± 1.7%), 9-octadecenoic acid (18:1n9; 24.3 ± 0.6%), octadecanoic acid (18:0; 20.0 ± 0.1%), 9,12-octadecadienoic acid (18:2n6; 8.6 ± 0.1%), 11-octadecenoic acid (18:1n7; 5.5 ± 0.5%), tetradecanoic acid (14:0; 4.7 ± 0.8%) and trace amounts (less than 1%) of heptadecanoic acid, 9-hexadecenoic acid, pentadecanoic acid, 15-methyl-hexadecanoic acid, 14-methyl-hexadecanoic acid, 10-heptadecenoic acid, 13-methyl-tetradecanoic acid and 12-methyl-tetradecanoic acid (Table 2).

### Genome properties

A phylogenetic tree highlighting the position of *Clostridium phoceensis* GD3<sup>T</sup> relative to other type strains within the order *Clostridiales* is provided in Fig. 4.

The genome of the *C. phoceensis* strain GD3<sup>T</sup> is 3 453 562 bp long with 59.32% G+C content. A total of 3320 genes were predicted, of which 3264 were protein-coding




FIG. 3. Electron microscopy of *Clostridium phoceensis* strain GD3<sup>T</sup>.

TABLE 1. Classification, general features and biochemical tests of *Clostridium phoceensis* strain GD3<sup>T</sup>


Property	Term
Current classification	Domain: <i>Bacteria</i> Phylum: <i>Firmicutes</i> Class: <i>Clostridia</i> Order: <i>Clostridiales</i> Family: <i>Clostridiaceae</i> Genus: <i>Clostridium</i> Species: <i>Clostridium phoceensis</i> Type strain: GD3
Gram stain	Negative
Cell shape	Bacillus
Cell diameter (µm)	0.5 µm
Cell length	1.8 µm
Motility	No
Sporulation	Yes
Temperature range	Mesophilic
Production of:	
Alkaline phosphatase	No
Catalase	No
Oxidase	No
Nitrate reductase	No
Urease	No
β-Galactosidase	No
N-acetyl-glucosamine	No
Esterase	Yes
Acid from:	
L-Arabinose	No
Ribose	No
Mannose	No
Mannitol	No
Sucrose	No
D-Glucose	Yes
D-Fructose	No
D-Maltose	No
D-Lactose	No
Habitat	Human

TABLE 2. Cellular fatty acid composition (%) of *Clostridium phoceensis* strain GD3<sup>T</sup>

Fatty acid	Name	Mean relative % <sup>a</sup>
16:0	Hexadecanoic acid	33.1 ± 1.7
18:1n9	9-Octadecenoic acid	24.3 ± 0.6
18:0	Octadecanoic acid	20.0 ± 0.1
18:2n6	9,12-Octadecadienoic acid	8.6 ± 0.1
18:1n7	11-Octadecenoic acid	5.5 ± 0.5
14:0	Tetradecanoic acid	4.7 ± 0.8
17:0	Heptadecanoic acid	TR
16:1n7	9-Hexadecenoic acid	TR
15:0	Pentadecanoic acid	TR
17:0 iso	15-methyl-Hexadecanoic acid	TR
17:0 anteiso	14-methyl-Hexadecanoic acid	TR
17:1n7	10-Heptadecenoic acid	TR
15:0 iso	13-Methyl-tetradecanoic acid	TR
15:0 anteiso	12-Methyl-tetradecanoic acid	TR

genes and 56 were RNA genes (three genes are 5S rRNA, one gene is 16S rRNA, one gene is 23S rRNA and 51 genes are tRNA genes). A total of 1967 genes (60.26%) were assigned a putative function. A total of 227 genes (6.95%) were identified as ORFans. The remaining genes (28.92%) were annotated as hypothetical proteins. The properties and statistics of the genome are summarized in Table 3 and Fig. 5. The distribution of genes into COGs functional categories is presented in Table 4 and Fig. 6.

**FIG. 4.** Phylogenetic tree highlighting position of *Clostridium phoceensis* GD3<sup>T</sup> relative to other type strains within *Clostridiales* order. GenBank accession numbers are indicated. Sequences were aligned using CLUSTALW, and phylogenetic inferences were obtained using maximum-likelihood method within MEGA 4 software [23]. Numbers at nodes are bootstrap values obtained by repeating analysis 500 times to generate majority consensus tree. Scale bar represents 5% nucleotide sequence divergence. *Rubidus massiliensis* was used as outgroup.



**Genome comparison**

We made some brief comparisons against nine genomes: *Intestinimonas butyriciproducens* strain ER1 (GenBank accession no. JPD01000000), *Flavonifractor plautii* ATCC 29863 (AGCK01000000), *Clostridium leptum* DSM 753 (ABCB02000000), *Clostridium cellulosi* DG5 (NZ\_LM995447), *Ethanoligenens harbinense* YUAN-3 (NC\_01482), *Oscillibacter valericigenes* Sjm18-20 (NC\_016048), *Clostridium viride* DSM 6836 (NZ\_JHZO01000000), *Eubacterium siraeum* V10Sc8a (FP929059) and *Pseudoflavonifractor capillosus* ATCC 29799 (NZ\_AAXG02000000). The draft genome sequence of *C. phoceensis* is smaller than those of *Pseudoflavonifractor*


*capillosus*, *Oscillibacter valericigenes*, *Flavonifractor plautii*, *Clostridium cellulosi* and *Intestinimonas butyriciproducens* (3.45, 4.24, 4.47, 3.81, 5.68 and 3.57 MB respectively), but larger than those of *Clostridium viride*, *Ethanoligenens harbinense*, *Clostridium leptum* and *Eubacterium siraeum* (2.41, 3.01, 2.82 and 2.84 MB respectively). The G+C content of *C. phoceensis* is smaller than those of *Flavonifractor plautii* (59.32 and 61.07% respectively) but larger than those of *Pseudoflavonifractor capillosus*, *Oscillibacter valericigenes*, *Clostridium viride*, *Ethanoligenens harbinense*, *Clostridium leptum*, *Eubacterium siraeum*, *Clostridium cellulosi* and *Intestinimonas butyriciproducens* (59.11, 53.19, 49.28, 55.56, 50.25, 45.13, 42.07 and 58.44% respectively). The gene content of *C. phoceensis* is smaller than those of *Pseudoflavonifractor capillosus*, *Oscillibacter valericigenes*, *Flavonifractor plautii*, *Clostridium cellulosi* and *Intestinimonas butyriciproducens* (3264, 4829, 4723, 4278, 5171 and 3529 respectively) but larger than those of *Clostridium viride*, *Ethanoligenens harbinense*, *Clostridium leptum* and *Eubacterium siraeum* (2321, 2701, 2482 and 2211 respectively). *C. phoceensis* has a similar distribution of genes into COGs categories with the most of the compared species (Fig. 4). However, *Clostridium cellulosi* was overrepresented for all the categories, and *Clostridium viride* was overrepresented for category Z (cytoskeleton).

**TABLE 3.** Nucleotide content and gene count levels of genome

Attribute	Genome (total)	
	Value	% of total <sup>a</sup>
Genome size (bp)	3 453 562	100
DNA coding region (bp)	2 924 785	84.68
DNA G+C content (bp)	2 044 366	59.31
Total genes	3320	100
rRNA	5	0.136
tRNA	51	0.116
Protein-coding genes	3264	98.31
Genes with function prediction	3111	17.02
Genes assigned to COGs	2534	57.39
Pseudo genes	62	1.86
Genes in internal clusters	1060	31.92
Genes with Pfam domains	2817	84
Genes with signal peptides	414	12.68
Genes with transmembrane helices	722	22.12
ORFan genes	227	6.95
CRISPR repeats	14	0.02

COGs, Clusters of Orthologous Groups database; CRISPR, clustered regularly interspaced short palindromic repeat.  
<sup>a</sup>Total is based on either size of genome in base pairs or total number of protein-coding genes in annotated genome.

In addition, *Clostridium phoceensis* shared 846, 884, 492, 562, 482, 717, 634, 407 and 842 orthologous genes respectively with *Intestinimonas butyriciproducens*, *Flavonifractor plautii*, *Clostridium leptum*, *Clostridium cellulosi*, *Ethanoligenens harbinense*, *Oscillibacter valericigenes*, *Clostridium viride*, *Eubacterium siraeum* and *Pseudoflavonifractor capillosus* (Table 5). Of these species, the orthologous genes shared ranged from 365 between *Eubacte-*



**FIG. 5.** Graphical circular map of *Clostridium phoceensis* GD3<sup>T</sup> genome. From outside in, outer two circles show ORFs oriented in forward (colored by COGs categories) and reverse (colored by COGs categories) directions, respectively. Third circle marks tRNA genes (green). Fourth circle shows G+C% content plot. Innermost circle shows GC skew, with purple indicating negative values and olive indicating positive values. COGs, Clusters of Orthologous Groups database; ORF, open reading frame.

**TABLE 4.** Number of genes associated with 25 general COGs functional categories

Code	Description	Value	% of total <sup>a</sup>
J	Translation	150	4.59
A	RNA processing and modification	0	0
K	Transcription	212	6.49
L	Replication, recombination and repair	106	3.24
B	Chromatin structure and dynamics	0	0
D	Cell cycle control, mitosis and meiosis	22	0.67
Y	Nuclear structure	0	0
V	Defense mechanisms	75	2.29
T	Signal transduction mechanisms	94	2.87
M	Cell wall/membrane biogenesis	54	1.65
N	Cell motility	27	0.82
Z	Cytoskeleton	0	0
W	Extracellular structures	0	0
U	Intracellular trafficking and secretion	25	0.76
O	Posttranslational modification, protein turnover, chaperones	51	1.56
C	Energy production and conversion	94	2.87
G	Carbohydrate transport and metabolism	106	3.24
E	Amino acid transport and metabolism	217	6.54
F	Nucleotide transport and metabolism	49	1.50
H	Coenzyme transport and metabolism	63	1.93
I	Lipid transport and metabolism	53	1.62
P	Inorganic ion transport and metabolism	94	2.87
Q	Secondary metabolites biosynthesis, transport and catabolism	29	0.88
R	General function prediction only	231	7.07
S	Function unknown	126	3.86
—	Not in COGs	1684	51.59

COGs, Clusters of Orthologous Groups database.

<sup>a</sup>Total is based on total number of protein-coding genes in annotated genome.

*rium siraeum* and *Clostridium viride* to 1079 between *Flavonifractor plautii* and *Intestinimonas butyriciproducens*. Compared to other species, *C. phoceensis* exhibited AGIOS values ranging from 56.64 with *Ethanoligenens harbinense* to 71.58 with *Flavonifractor plautii*.

## Conclusion


On the basis of taxonogenomic analyses, we propose *Clostridium phoceensis* sp. nov., strain GD3<sup>T</sup> (= CSUR P1929 = DSM 100334), as the type strain of *C. phoceensis* sp. nov., a new species within the genus *Clostridium* (Fig. 4). This strain was isolated from the gut microbiota of a 28-year-old healthy French man.

## Taxonomic and nomenclatural proposals

### Description of *Clostridium phoceensis* sp. nov.

*Clostridium phoceensis* strain GD3<sup>T</sup> (pho.ce.en'sis, L. fem. adj., from *phoceensis* referring to Phocea, the Greek name of the city which founded Marseille, where it was isolated) is a Gram-negative nonmotile bacillus whose individual cell size is

**FIG. 6.** Distribution of functional classes of predicted genes in genomes from *Clostridium phoceensis* (Cp), *Intestinimonas butyriciproducens* strain ER1 (Ib), *Flavonifractor plautii* ATCC 29863 (Fp), *Clostridium leptum* DSM 753 (Cl), *Clostridium cellulosi* DG5 (Cc), *Ethanoligenens harbinense* YUAN-3 (Eh), *Oscillibacter valericigenes* Sjm18-20 (Ov), *Clostridium viride* DSM 6836 (Cv), *Eubacterium siraeum* V10Sc8a (Es) and *Pseudoflavonifractor capillosus* ATCC 29799 (PC) genomes according to clusters of orthologous groups of proteins. ATCC, American Type Culture Collection (Manassas, VA, USA).



**TABLE 5.** Orthologous genes shared (upper right) and AGIOS values obtained (lower left)<sup>a</sup>

	Cp	Ib	Fp	Cl	Cc	Eh	Ov	Cv	Es	Pc
Cp	<b>3264</b>	846	884	492	562	482	717	634	407	842
Ib	70.10	<b>3529</b>	1079	508	627	501	771	671	407	1016
Fp	71.58	73.64	<b>4278</b>	523	649	518	818	686	433	1039
Cl	61.83	61.60	61.65	<b>2482</b>	587	495	475	432	405	523
Cc	57.17	57.91	56.82	61.62	<b>5171</b>	619	604	562	465	616
Eh	56.64	56.86	56.92	56.13	55.34	<b>2701</b>	515	449	415	499
Ov	65.72	65.34	65.55	61.60	58.40	56.54	<b>4723</b>	653	409	710
Cv	63.84	64.57	64.21	60.85	59.13	55.02	62.58	<b>2321</b>	364	614
Es	57.83	58.77	57.72	60.78	61.38	54.37	59.13	58.93	<b>2211</b>	427
Pc	61.55	61.95	62.89	55.43	52.37	57.21	58.02	56.73	53.24	<b>4829</b>

AGIOS, average genomic identity of orthologous gene sequences; ATCC, American Type Culture Collection (Manassas, VA, USA).

<sup>a</sup>Values in bold are gene numbers. Ten genomes were used for this study: *Clostridium phoceensis* (Cp), *Intestinimonas butyriciproducens* strain ER1 (Ib), *Flavonifractor plautii* ATCC 29863 (Fp), *Clostridium leptum* DSM 753 (Cl), *Clostridium cellulosi* DG5 (Cc), *Ethanoligenens harbinense* YUAN-3 (Eh), *Oscillibacter valericigenes* Sjm18-20 (Ov), *Clostridium viride* DSM 6836 (Cv), *Eubacterium siraeum* V10Sc8a (Es) and *Pseudoflavonifractor capillosus* ATCC 29799 (PC) genomes according to the clusters of orthologous groups of proteins.

1.8 µm in length and 0.5 µm in diameter. It is a strictly anaerobic and endospore-forming bacterium. Strain GD3<sup>T</sup> is catalase and oxidase negative, and its optimal growth temperature is 45°C, but it also grows weakly at 37°C. Biochemical analyses showed positive reactions of *C. phoceensis* for D-glucose and produced esterase (C4), phosphatase acid, naphthol-AS-BI-phosphohydrolase and β-glucosidase enzymes. *C. phoceensis* was sensitive to amoxicillin, amoxicillin-clavulanate, ceftriaxone, ceftazidime, imipenem, doripenem and ciprofloxacin. Its predominant cellular fatty acids are hexadecanoic acid (16:0; 33.1 ± 1.7%), 9-octadecenoic acid (18:1n9; 24.3 ± 0.6%), octadecanoic acid (18:0; 20.0 ± 0.1%), 9,12-octadecadienoic acid (18:2n6; 8.6 ± 0.1%), 11-octadecenoic acid (18:1n7; 5.5 ± 0.5%) and tetradecanoic acid (14:0; 4.7 ± 0.8%). Its 16S rRNA sequences were deposited in GenBank under accession numbers [CVUG01000000](#) and [LN846907](#). Strain GD3<sup>T</sup>, whose CSUR

and DSMZ numbers are respectively CSUR P1929 and DSM 100334, was identified from the stool sample of a 28-year-old healthy French man.

### Acknowledgements

The authors thank the Xegen Company ([www.xegen.fr](http://www.xegen.fr)) for automating the genomic analysis process and culturomics team in URMITE for the bacteria isolation. This project was funded by the Fondation Méditerranée Infection.

### Conflict of Interest

None declared.

## References

- [1] Palmer C, Bik EM, DiGiulio DB, Relman DA, Brown PO. Development of the human infant intestinal microbiota. *PLoS Biol* 2007;5:e177.
- [2] Lagier JC, Million M, Hugon P, Armougom F, Raoult D. Human gut microbiota: repertoire and variations. *Front Cell Infect Microbiol* 2012;2:136.
- [3] Lagier JC, Edouard S, Pagnier I, Mediannikov O, Drancourt M, Raoult D. Current and past strategies for bacterial culture in clinical microbiology. *Clin Microbiol Rev* 2015;28:208–36.
- [4] Ramasamy D, Mishra AK, Lagier JC, Padhmanabhan R, Rossi M, Sentausa E, et al. A polyphasic strategy incorporating genomic data for the taxonomic description of novel bacterial species. *Int J Syst Evol Microbiol* 2014;64:384–91.
- [5] Lawson PA, Rainey FA. Proposal to restrict the genus *Clostridium* (Prazmowski) to *Clostridium butyricum* and related species. *Int J Syst Evol Microbiol* 2015.
- [6] Popoff MR, Bouvet P. Clostridial toxins. *Future Microbiol* 2009;4:1021–64.
- [7] Lagier JC, Armougom F, Million M, Hugon P, Pagnier I, Robert C, et al. Microbial culturomics: paradigm shift in the human gut microbiome study. *Clin Microbiol Infect* 2012;18:1185–93.
- [8] Grosse-Herrenthey A, Maier T, Gessler F, Schaumann R, Böhnelt H, Kostrzewa M, et al. Challenging the problem of clostridial identification with matrix-assisted laser desorption and ionization–time-of-flight mass spectrometry (MALDI-TOF MS). *Anaerobe* 2008;14:242–9.
- [9] Seng P, Abat C, Rolain JM, Colson P, Lagier JC, Gouriet F, et al. Identification of rare pathogenic bacteria in a clinical microbiology laboratory: impact of matrix-assisted laser desorption ionization–time of flight mass spectrometry. *J Clin Microbiol* 2013;51:2182–94.
- [10] Drancourt M, Bollet C, Carlioz A, Martelin R, Gayral JP, Raoult D. 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacterial isolates. *J Clin Microbiol* 2000;38:3623–30.
- [11] Dione N, Sankar SA, Lagier JC, Khelaifia S, Michele C, Armstrong N, et al. Genome sequence and description of *Anaerosalibacter massiliensis* sp. nov. *New Microbes New Infect* 2016;10:66–76.
- [12] Mourembou G, Yasir M, Azhar El, Lagier JC, Bibi F, Jiman-Fatani AA, et al. Rise of microbial culturomics: noncontiguous finished genome sequence and description of *Beduini massiliensis* gen. nov., sp. nov. *OMICS J Integr Biol* 2015;19:766–76.
- [13] Fournier PE, Raoult D, Drancourt M. New species announcement: a new format to prompt the description of new human microbial species. *New Microbes New Infect* 2016. In press.
- [14] Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol* 2008;26:541–7.
- [15] Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 2010;11:119.
- [16] Benson DA, Karsch-Mizrachi I, Clark K, Lipman DJ, Ostell J, Sayers EW. GenBank. *Nucleic Acids Res* 2012;40(Database issue):D48–53.
- [17] Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 1997;25:955–64.
- [18] Lagesen K, Hallin P, Rødland EA, Stærfeldt HH, Rognes T, Ussery DW. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 2007;35:3100–8.
- [19] Dyrlov Bendtsen J, Nielsen H, von Heijne G, Brunak S. Improved prediction of signal peptides: SignalP 3.0. *J Mol Biol* 2004;340:783–95.
- [20] Krogh A, Larsson B, von Heijne G, Sonnhammer ELL. Predicting transmembrane protein topology with a hidden markov model: application to complete genomes. *J Mol Biol* 2001;305:567–80.
- [21] Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, et al. Artemis: sequence visualization and annotation. *Bioinformatics* 2000;16:944–5.
- [22] Carver T, Thomson N, Bleasby A, Berriman M, Parkhill J. DNAPlotter: circular and linear interactive genome visualization. *Bioinformatics* 2009;25:119–20.
- [23] Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol* 2007;24:1596–9.